

Mikael Elias,^a Jérôme Dupuy,^b
Luigia Merone,^c Claude
Lecomte,^a Mosè Rossi,^c Patrick
Masson,^d Giuseppe Manco^c and
Eric Chabriere^{a,d*}

^aLaboratoire de Cristallographie et Modélisation des Matériaux Minéraux et Biologiques, CNRS–Université Henri Poincaré, 54506 Vandœuvre-lès-Nancy, France, ^bLaboratoire de Cristallogénèse et Cristallographie des Protéines, Institut de Biologie Structurale J.-P. Ebel, 38027 Grenoble, France, ^cIstituto di Biochimica delle Proteine, Consiglio Nazionale delle Ricerche, Via P. Castellino 111, 80131 Napoli, Italy, and ^dUnité d'Enzymologie, Département de Toxicologie, Centre de Recherches du Service de Santé des Armées, 38702 La Tronche, France

Correspondence e-mail:
eric.chabriere@lcm3b.uhp-nancy.fr

Received 12 March 2007
Accepted 13 May 2007



© 2007 International Union of Crystallography
All rights reserved

Crystallization and preliminary X-ray diffraction analysis of the hyperthermophilic *Sulfolobus solfataricus* phosphotriesterase

Organophosphates constitute the largest class of insecticides used worldwide and some of them are potent nerve agents. Consequently, organophosphate-degrading enzymes are of paramount interest as they could be used as bioscavengers and biodecontaminants. Phosphotriesterases (PTEs) are capable of hydrolyzing these toxic compounds with high efficiency. A distant and hyperthermophilic representative of the PTE family was cloned from the archeon *Sulfolobus solfataricus* MT4, overexpressed in *Escherichia coli* and crystallized; the crystals diffracted to 2.54 Å resolution. Owing to its exceptional thermostability, this PTE may be an excellent candidate for obtaining an efficient organophosphate biodecontaminant. Here, the crystallization conditions and data collection for the hyperthermophilic *S. solfataricus* PTE are reported.

1. Introduction

Organophosphates (OPs) are toxic compounds because they irreversibly inhibit acetylcholinesterase, a key enzyme in the nervous system. They have been distributed globally since the end of World War II and their toxic properties have been utilized in chemical warfare agents such as sarin, soman and VX and for the production of agricultural insecticides (Raushel, 2002). Enzymes that are capable of degrading OPs are currently extensively studied because of their potential in decontamination and detection systems for organophosphate-based pesticides and nerve agents (LeJeune *et al.*, 1998). Enzymatic detoxification of OPs has become the subject of numerous studies as current methods such as bleach treatment and incineration are impractical owing to high costs or environmental concerns. Bacterial OP hydrolases are appealing for this application owing to their broad substrate specificity and high catalytic rate under mild conditions (LeJeune *et al.*, 1998).

Enzymes that catalyse the hydrolysis of phosphoester bonds in OPs are known from several different organisms. One class of enzymes that are able to degrade nerve gases is the prolidases. As an example, a prolidase named organophosphorus acid anhydrolase (OPAA) was identified in a strain of *Alteromonas* (Cheng *et al.*, 1993). Another class of enzyme, isolated from *Loligo vulgaris*, is diisopropylfluorophosphatase (DFPase), which prefers P–F bonds. Its three-dimensional structure was solved and reveals a six-bladed β -propeller with two calcium ions in a central water-filled tunnel (Koepke *et al.*, 2002). HDL-associated human paraoxonase (HPON) also hydrolyses phosphotriesters, albeit with lower proficiency. The structure of HPON1 has recently been solved and shows a similar fold to that of DFPase (Harel *et al.*, 2004). Other bacterial enzymes (EC 3.1.8.1) called phosphotriesterases (PTE) or sometimes organophosphorus hydrolases (OPH), organophosphate-degrading enzymes (OPD), parathion hydrolases (Hou *et al.*, 1996) or paraoxonases (Merone *et al.*, 2005), show a preference for organophosphorus compounds containing P–O or P–S bonds. Phosphotriesterases are members of the the amidohydrolase superfamily (Seibert & Raushel, 2005), enzymes that catalyze the hydrolysis of a wide range of compounds with different chemical properties (phosphoesters, esters, amides *etc.*). Several different structures of PTEs are available. Structures are known of the most efficient PTE, that from *Pseudo-*

monas diminuta (Munnecke, 1976), and of the very similar (90% sequence identity) OpdA from *Agrobacterium radiobacter* (Horne *et al.*, 2003). Structurally, PTEs are $(\beta/\alpha)_8$ -barrel enzymes with binuclear metal centres located at the C-terminal end of the barrel (Vanhook *et al.*, 1996). The reaction mechanism was proposed to proceed via an S_N2 -like mechanism in which the metal centre enables a hydroxide ion (bridged to the two metal ions) to attack the electrophilic phosphorus of the substrate (Aubert *et al.*, 2004).

A protein from the hyperthermophilic archeon *Sulfolobus solfataricus* MT4, *SsoPox*, has recently been cloned and characterized: it is a rather distant representative of the PTE family, displaying only about 30% sequence identity to mesophilic PTEs. Despite this, all the amino acids coordinating the binuclear metal centre are conserved. It has been observed that *SsoPox* catalyzes the hydrolysis of paraoxon and other pesticides with a significantly lower proficiency than mesophilic PTEs, but with a similar K_m . Similarly to *P. diminuta* PTE, its activity depends on the presence of metal cations; high activity was observed for the Co^{2+} -substituted isoenzyme (Merone *et al.*, 2005). *SsoPox* proved to have exceptional thermal stability, with denaturation half-lives of 4 h and 90 min at 368 and 373 K, respectively. This property allows high-yield purification of recombinant enzyme simply by heating cell lysates to cause precipitation of host (*Escherichia coli*) proteins. Recently, high catalytic activity and specificity towards lactones as substrates was reported for *SsoPox* and other PTE-related enzymes. Afriat *et al.* (2006) dubbed this new group of enzymes 'phosphotriesterases like lactonases' (PLLs) based on the observation of sequence features that were not present in mesophilic PTEs and of significant differences in enzyme specificity.

In particular, the activity detected towards natural homoserine lactones suggests a role in quorum-sensing signalling and led to the conclusion that PTEs evolved from a PLL-family member, probably *SsoPox*, utilizing its latent promiscuous phosphotriesterase activity as an essential starting point (Afriat *et al.*, 2006). Furthermore, although its phosphotriesterase activity at ambient temperature against OPs is very low, the structure of *SsoPox* may provide clues to enhancing its catalytic parameters. Finally, *SsoPox* may be an excellent candidate in biotechnology studies seeking an efficient biodecontaminant of organophosphorus compounds.

In this report, we describe the crystallization, data collection and preliminary X-ray diffraction analysis of the hyperthermophilic *S. solfataricus* phosphotriesterase.




Figure 1
Typical crystal of the hyperthermophilic phosphotriesterase from *S. solfataricus*.

2. Crystallization

For the crystallization studies presented here, we utilized recombinant *SsoPox* overproduced in *E. coli* under the control of the T7 promoter and extensively purified in a soluble and active form (Merone *et al.*, 2005). Crystallization assays were carried out using the hanging-drop vapour-diffusion technique (McPherson, 1990). The protein solution for crystallization was made up of 20 mM HEPES pH 8.5, 0.2 mM $CoCl_2$ and 0.2 M NaCl. The enzyme was concentrated to 5.8 mg ml⁻¹ using a centrifugation device (Centriprep Amicon, 10 kDa cutoff; Millipore, St Quentin-en-Yvelines, France). Equal volumes (1–2 μ l) of protein and reservoir solutions were mixed and the resulting drops were equilibrated against 800 μ l reservoir solution containing 15–18% (w/v) PEG 8000 and 50 mM Tris–HCl buffer pH 8. Small crystals appeared after one week at 277 K (Fig. 1). Although they did not possess clear and geometrical faces, the crystals diffracted to reasonable resolution (2.54 Å).

3. Data collection

Crystals were mounted in nylon loops (Hampton Research) and flash-frozen in liquid nitrogen at 100 K using a cryoprotectant solution containing 50 mM Tris–HCl buffer pH 8, 18% (w/v) PEG 8000 and 25% (v/v) glycerol. X-ray diffraction intensities were collected at the BM30A beamline (ESRF, Grenoble, France) using a wavelength of 0.9789 Å and a MAR CCD 165 mm detector (MAR Research) with 35 s exposures. Diffraction data were collected from 180 images using the oscillation method; individual frames consisted of 1.0° oscillation steps over a range of 180° (Fig. 2).

4. Results and conclusion

Crystals of *SsoPox* belong to the orthorhombic space group $P2_12_12_1$, with unit-cell parameters $a = 87.16$, $b = 104.82$, $c = 155.36$ Å (Table 1). X-ray diffraction data were integrated, scaled and merged using the




Figure 2
A diffraction pattern of a crystal of the phosphotriesterase from *S. solfataricus*. The edge of the frame is at 2.1 Å.

Table 1

Diffraction data collected at BM30A FIP beamline (ESRF, Grenoble, France).

Values in parentheses are for the highest resolution shell.

| | |
|-----------------------------|-------------------------------------|
| Wavelength (Å) | 0.9789 |
| Resolution (Å) | 2.54 (2.7–2.54) |
| Space group | $P2_12_12_1$ |
| Unit-cell parameters (Å) | $a = 87.16, b = 104.82, c = 155.36$ |
| No. of observed reflections | 308664 (50951) |
| No. of unique reflections | 85611 (12853) |
| Completeness (%) | 95.3 (87.2) |
| R_{merge} (%) | 16.0 (52.8) |
| R_{meas} (%) | 11.6 (52) |
| $I/\sigma(I)$ | 11.69 (3.03) |
| Redundancy | 3.60 (3.96) |
| Completeness (%) | 95.3 (87.2) |

XDS program (Kabsch, 1993) and the *CCP4* program suite (Collaborative Computational Project, Number 4, 1994). Initial molecular replacement was performed with *Phaser* (Read, 2001) using a poly-alanine model derived from the structure of *P. diminuta* PTE (PDB code 1dpm; Vanhooke *et al.*, 1996). Two protein molecules were found in the asymmetric unit (translation-function *Z* scores of 7.22 and 6.72). Given the molecular weight of *SsoPox* (35.5 kDa), this corresponds to a high Matthews coefficient (V_M) of $5.0 \text{ \AA}^3 \text{ Da}^{-1}$ and a high solvent content of approximately 75.2%. The electron-density map calculated with model phases obtained from molecular replacement was of poor quality. Almost all side chains were not visible in the maps. Nevertheless, the active site constituted of two metal cations was clearly visible. The electron-density map was improved with *DM* (Cowtan & Zhang, 1999) using solvent flattening and noncrystallographic symmetry averaging between the two molecules (starting FOM, 0.442; FOM after *DM*, 0.632). Manual model improvement was performed using *Coot* (Emsley & Cowtan, 2004). Some side chains could be correctly placed and some loops involved in the dimer interface were removed. Using the improved model and fixing the two initial solutions, molecular replacement was performed using *MOLREP* (Vagin & Teplyakov, 2000) and two extra solutions emerged, giving a total of four protein molecules in the asymmetric unit ($R_{\text{cryst}} = 0.47$, correlation coefficient = 0.325). This corresponds to a V_M of $2.5 \text{ \AA}^3 \text{ Da}^{-1}$, which is a more typical value (Matthews, 1968).

Finally, the new electron-density maps were of sufficient quality for model building. Structure refinement and interpretation are in progress.

This research was supported by grants to PM and EC from Délégation Générale pour l'Armement (CO No. 010807/03-10) and from the CNRS. We also thank MIUR project 'Piano Nazionale Ricerca per le Biotecnologie Avanzate II–Biocatalisi'.

References

- Afriati, L., Roodvelt, C., Manco, G. & Tawfik, D. (2006). *Biochemistry*, **45**, 13677–13686.
- Aubert, S. D., Li, Y. & Raushel, F. M. (2004). *Biochemistry*, **43**, 5707–5715.
- Cheng, T. C., Harvey, S. P. & Stroup, A. N. (1993). *Appl. Environ. Microbiol.* **59**, 3138–3140.
- Collaborative Computational Project, Number 4 (1994). *Acta Cryst.* **D50**, 760–763.
- Cowtan, K. D. & Zhang, K. Y. (1999). *Prog. Biophys. Mol. Biol.* **72**, 245–270.
- Emsley, P. & Cowtan, K. (2004). *Acta Cryst.* **D60**, 2126–2132.
- Harel, M., Aharoni, A., Gaidukov, L., Brumshtein, B., Khersonsky, O., Meged, R., Dvir, H., Ravelli, R. B., McCarthy, A., Tokor, L., Silman, I., Sussman, J. L. & Tawfik, D. S. (2004). *Nature Struct. Mol. Biol.* **11**, 412–419.
- Hou, X., Maser, R. L., Magenheimer, B. S. & Calvet, J. P. (1996). *Gene*, **168**, 157–163.
- Horne, I., Qiu, X., Russel, R. J. & Oakeshott, J. G. (2003). *FEMS Microbiol. Lett.* **222**, 1–8.
- Kabsch, W. (1993). *J. Appl. Cryst.* **26**, 795–800.
- Koepke, J., Scharff, E. I., Lucke, C., Ruterjans, H. & Fritzsche, G. (2002). *Acta Cryst.* **D58**, 1757–1759.
- LeJeune, K. E., Wild, J. R. & Russell, A. J. (1998). *Nature (London)*, **395**, 27–28.
- McPherson, A. (1990). *Eur. J. Biochem.* **189**, 1–23.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- Merone, L., Mandrich, L., Rossi, M. & Manco, G. (2005). *Extremophiles*, **9**, 297–305.
- Munnecke, D. M. (1976). *Appl. Environ. Microbiol.* **32**, 7–13.
- Raushel, F. M. (2002). *Curr. Opin. Microbiol.* **5**, 288–295.
- Read, R. J. (2001). *Acta Cryst.* **D57**, 1373–1382.
- Seibert, C. M. & Raushel, F. M. (2005). *Biochemistry*, **44**, 6383–6391.
- Vagin, A. & Teplyakov, A. (2000). *Acta Cryst.* **D56**, 1622–1624.
- Vanhooke, J. L., Benning, M. M., Raushel, F. M. & Holden, H. M. (1996). *Biochemistry*, **35**, 6020–6025.